

Surgical Threshold Policy

Knee Arthroscopy for Chronic Knee Pain

Scope

This policy covers the use of knee arthroscopy as a surgical intervention or diagnostic tool for chronic knee pain in adults. It does not cover arthroscopy recommended by an orthopaedic specialist in those under 18 years of age or in adults following acute injury (see Note 1) with suspected internal joint derangement, septic arthritis or suspected malignancy. Autologous chondrocyte implantation (ACI) is funded by NHS England in specialist centres as per criteria in [NICE TA508](#) (see reference 1).

Policy

Referral for treatment should be through the MSK service/pathway.

It is the responsibility of referring and treating clinicians to ensure compliance with this policy. Referral proforma should be attached to the patient notes to aid the clinical audit process and provide evidence of compliance with the policy. For patients not meeting the policy criteria, clinicians can apply for funding to the Exceptional Cases Panel by completing the exceptional funding section of the referral proforma:

Primary/Community Care

Patients should be referred as per the pathway overleaf.

In patients with chronic knee pain, where evidence is provided that a comprehensive course of conservative treatment (rest, lifestyle advice, optimum pharmacological treatments and physiotherapy) has been tried for 12 months and failed, patients can be referred by physiotherapy/MSK for specialist consultation.

Where patients have non-osteoarthritis symptoms that worsen with conservative treatment, physiotherapy/MSK may choose to make a referral after shorter periods of conservative treatment.

Referral for MRI scans should only be made by secondary care consultants or specialists working in NHS C&P (ICB) commissioned MSK services.

Secondary Care

Where a competent clinical examination or MRI scan has demonstrated clear evidence of a significant internal joint derangement which is likely to be the cause of the symptoms and suitable for treatment by knee arthroscopy, surgery will be funded.

However, the following are **Lower Clinical Priority** procedures and will not be funded without Exceptional Cases Panel approval:

1. Lavage or debridement of osteoarthritic knees, except in cases of true mechanical locking (see note 2).
2. Repair or debridement of degenerative meniscal tears, except in cases of true mechanical locking.
3. Treatment of Patellofemoral pain syndrome, patellar tendinopathy or patellar dislocation except in cases where there is presence of loose bodies.

4. Use as a diagnostic tool, except in cases where there is on-going diagnostic uncertainty following both clinical examination and non-invasive imaging procedures (e.g. MRI) conducted by specialists.

Hyaluronic acid injections: Intra-articular or periarticular hyaluronic acid or its derivatives is not routinely commissioned for injection into the peripheral joints of the knee, hips, ankle or thumb.

Smoking

Patients who smoke should be advised to attempt to stop smoking - see [stop smoking policy](#).


Weight Management

Patients who are overweight or obese should be offered referral to the appropriate weight management service.

Notes

- Note 1:** Suspected acute/traumatic onset meniscus tear should have a significant cause (e.g. sport, twisting while running, external force on the knee), compared to gradual onset that would be more consistent with degenerative meniscus tear.
- Note 2:** An intermittent block to normal range of movement (usually extension). These patients should be referred urgently.

Pathway



*Details available at: <http://www.eoemskservice.nhs.uk/advice-and-exercises>

Procedures that are not routinely funded

For some interventions, the evidence identifies a lack of effect and insufficient evidence to warrant their use. RCTs show limited benefit of knee arthroscopy for patients with degenerative meniscal tears (repair, debridement) (see references 2,3,4 & 5). RCTs of patients with osteoarthritis of the knees also show no benefit of lavage and/or debridement (see references 3, 6 & 7). NICE guidance states that 'referral for arthroscopic lavage and debridement should not be offered as part of treatment for osteoarthritis, unless there is a clear history of mechanical locking (not gelling, 'giving way' or X-ray evidence of loose bodies).' Lavage in isolation is not indicated (see references 8, 9 & 10). In an RCT of patients with patellofemoral pain syndrome, mixed arthroscopic procedures and exercise resulted in equivalent improvements compared with exercise alone (see references 11 & 12). Similarly, trials of knee arthroscopy for patellar tendinopathy (see reference 13 & 14) and patellar dislocation (see reference 15 & 16) have not shown advantages over conservative management.

Other procedures

For arthroscopic articular cartilage surgery, including debridement of non-osteoarthritic joints, microfracture, autologous chondrocyte implantation and osteochondral autologous transplantation (Mosaicplasty), there is no robust evidence for their effectiveness (see reference 17). Although there is an absence of evidence for the benefit of these procedures, there is currently no evidence that they have no benefit. There is NICE guidance recommending the use of some procedures (see reference, 18, 19 & 20) but not others (see reference 21). Therefore, these procedures are not restricted as Lower Clinical Priority procedures, but patients should be treated conservatively, as per the pathway in this policy before they are considered for surgery.

Restricted use of MRI

MRI is a good diagnostic tool (see reference 22) but may be inaccurate when used by less experienced staff (see reference 23). It is not considered necessary for the majority of patients presenting in primary care and its use is, therefore, restricted to secondary care or MSK specialists for this indication.

Glossary

Arthroscopy:	Key-hole surgery using an arthroscope (light source and video camera).
Articular cartilage:	The cartilage covering the articular surfaces of the bones participating in a synovial joint.
Autologous chondrocyte implantation:	Harvesting of tissue from the edge of the affected knee joint, which is cultured and, in a second surgical procedure, implanted into damaged areas.
Chondroplasty:	Reparative surgery of cartilage.
Debridement:	Removal of dead, damaged, or infected tissue.
Lavage:	Cleaning or rinsing.

References

1. NICE.org.uk. Overview | Autologous chondrocyte implantation using chondrosphere for treating symptomatic articular cartilage defects of the knee | Guidance | NICE [Internet]. [cited 2023 Aug 10]. Available from: <https://www.nice.org.uk/guidance/ta508>
2. O'Connor D, Johnston R V., Brignardello-Petersen R, Poolman RW, Cyril S, Vandvik PO, et al. Arthroscopic surgery for degenerative knee disease (osteoarthritis including degenerative meniscal tears). Cochrane Database Syst Rev. 2022;2022(3).
3. Siemieniuk RAC, Harris IA, Agoritsas T, Poolman RW, Brignardello-Petersen R, Van de Velde S, et al. Arthroscopic surgery for degenerative knee arthritis and meniscal tears: a clinical practice guideline. BMJ. 2017 May;357:j1982.
4. Hohmann E, Glatt V, Tetsworth K, Cote M. Arthroscopic Partial Meniscectomy Versus Physical Therapy for Degenerative Meniscus Lesions: How Robust Is the Current Evidence? A Critical

Systematic Review and Qualitative Synthesis. *Arthrosc J Arthrosc Relat Surg Off Publ Arthrosc Assoc North Am Int Arthrosc Assoc.* 2018 Sep;34(9):2699–708.

5. Migliorini F, Oliva F, Eschweiler J, Cuozzo F, Hildebrand F, Maffulli N. No evidence in support of arthroscopic partial meniscectomy in adults with degenerative and nonobstructive meniscal symptoms: a level I evidence-based systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2023 May;31(5):1733–43.
6. Laupattarakasem W, Laopaiboon M, Laupattarakasem P, Sumananont C. Arthroscopic debridement for knee osteoarthritis. *Cochrane database Syst Rev.* 2008 Jan;(1):CD005118.
7. Reichenbach S, Rutjes AW, Nüesch E, Trelle S, Jüni P. Joint lavage for osteoarthritis of the knee. *Cochrane database Syst Rev.* 2010 May;(5):CD007320.
8. Ebi.aomrc.org.uk. Knee arthroscopy for patients with osteoarthritis - EBI [Internet]. [cited 2023 Aug 10]. Available from: <https://ebi.aomrc.org.uk/interventions/knee-arthroscopy-for-patients-with-osteoarthritis/>
9. NICE.org.uk. Osteoarthritis in over 16s: diagnosis and management [Internet]. NICE guideline. 2022. Available from: <https://www.nice.org.uk/guidance/ng226>
10. NICE.org.uk. Overview | Arthroscopic knee washout, with or without debridement, for the treatment of osteoarthritis | Guidance | NICE [Internet]. [cited 2023 Aug 9]. Available from: <https://www.nice.org.uk/guidance/ipg230>
11. Kettunen JA, Harilainen A, Sandelin J, Schlenzka D, Hietaniemi K, Seitsalo S, et al. Knee arthroscopy and exercise versus exercise only for chronic patellofemoral pain syndrome: A randomized controlled trial. *BMC Med.* 2007 Dec 13;5.
12. Kettunen JA, Harilainen A, Sandelin J, Schlenzka D, Hietaniemi K, Seitsalo S, et al. Knee arthroscopy and exercise versus exercise only for chronic patellofemoral pain syndrome: 5-year follow-up. *Br J Sports Med.* 2012 Mar;46(4):243–6.
13. Dan M, Phillips A, Johnston R V, Harris IA. Surgery for patellar tendinopathy (jumper's knee). *Cochrane Database Syst Rev.* 2019;(9).
14. Koëter S, Van Tienen TG, Rijk PC, Vincken PWJ, Segers MJM, Van Essen B, et al. Dutch Guideline on Knee Arthroscopy Part 2: non-menisus intra-articular knee injury: a multidisciplinary review by the Dutch Orthopaedic Association. *Acta Orthop.* 2021;92(1):81.
15. Xing X, Shi H, Feng S. Does surgical treatment produce better outcomes than conservative treatment for acute primary patellar dislocations? A meta-analysis of 10 randomized controlled trials. *J Orthop Surg Res.* 2020 Mar 24;15(1).
16. Smith TO, Gaukroger A, Metcalfe A, Hing CB. Surgical versus non-surgical interventions for treating patellar dislocation. *Cochrane database Syst Rev.* 2023 Jan 24;1(1).
17. Devitt BM, Bell SW, Webster KE, Feller JA, Whitehead TS. Surgical treatments of cartilage defects of the knee: Systematic review of randomised controlled trials. *Knee.* 2017 Jun 1;24(3):508–17.
18. NICE.org.uk. Overview | Mosaicplasty for symptomatic articular cartilage defects of the knee | Guidance | NICE [Internet]. [cited 2023 Aug 9]. Available from: <https://www.nice.org.uk/guidance/ipg607>
19. NICE.org.uk. 1 Recommendations | Autologous chondrocyte implantation for treating symptomatic articular cartilage defects of the knee | Guidance | NICE [Internet]. [cited 2023 Aug 9]. Available from: <https://www.nice.org.uk/guidance/ta477/chapter/1-Recommendations>
20. NICE.org.uk. Overview | Arthroscopic radiofrequency chondroplasty for discrete chondral defects of the knee | Guidance | NICE [Internet]. [cited 2023 Aug 9]. Available from: <https://www.nice.org.uk/guidance/ipg493>
21. NICE.org.uk. Overview | Microstructural scaffold (patch) insertion without autologous cell implantation for repairing symptomatic chondral knee defects | Guidance | NICE.
22. Crawford R, Walley G, Bridgman S, Maffulli N. Magnetic resonance imaging versus arthroscopy in the diagnosis of knee pathology, concentrating on meniscal lesions and ACL tears: a systematic

review. Br Med Bull. 2007;84(1):5–23.

23. Bryan S, Weatherburn G, Bungay H, Hatrick C, Salas C, Parry D, et al. The cost-effectiveness of magnetic resonance imaging for investigation of the knee joint. Health Technol Assess. 2001;5(27).

Policy effective from	Policy adopted by NHS C&P (ICB) 1 July 2022
Policy to be reviewed:	Reviewed policy ratified by QPF 22 March 2024
	Reviewed policy approved by JCPEG 13 March 2024
	Reviewed policy approved by CPF 5 March 2024
	Policy developed by C&P Public Health Team
Document Reference:	CCG - CP Forum and Policies - CPF Pols & Working Area\Policies, Pathways & Position Statements\3. Surgical Threshold Policies\Knee Arthroscopy\Policies